

PECAM1

Reactivity: Human

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW: 83kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human PECAM1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

PECAM1; CD31; FLJ58394; PECAM-1;

Catalog #: A2104

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 5175

Isotype: IgG

Swiss Prot: P16284

Purity: Affinity purification

For research use only.

Background:

CD31 (Platelet Endothelial Cell Adhesion Molecule-1: PECAM-1), a member of the Ig superfamily of cell adhesion molecules, is expressed by circulating platelets, monocytes, neutrophils, some T cells and endothelial cells, and modulates cell adhesion, endothelial cell migration and angiogenesis (1). CD31 is phosphorylated on Tyr686 at the cytoplasmic carboxy-terminal tail upon various stimuli (e.g. mechanical stress or oxidative stress), presumably by Src family members (2). The tyrosine phosphorylation mediates associations with a number of SH2 domain containing binding partners such as PI3-kinase, SHIP, PLC and SHP-2. Thus, CD31 serves as a scaffold for various signaling molecules (3).

To place an order, please [Click HERE](#).