PPP4C

Reactivity: Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200 Calculated MW:35kDa Observed MW:Refer to Figures Immunogen: Recombinant protein of human PPP4C Storage Buffer: Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3. Concentration:

e Synonym:

PP4; PPX; PP4C; PPH3; PPP4;

Background:

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified, designated PP1, PP2A, PP2B (calcineurin) and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4) is a putative member of a novel PP family.

To place an order, please Click HERE.



Catalog #:A2109 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:5531 Isotype:IgG Swiss Prot:P60510 Purity:Affinity purification

For research use only.



