

LCK

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:56kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human LCK

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

bd

Synonym:

YT16; p56lck; pp58lck;

Catalog #:A2177

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3932

Isotype:IgG

Swiss Prot:P06239

Purity:Affinity purification

For research use only.

Background:

The Src family of protein tyrosine kinases, which includes Src, Lyn, Fyn, Yes, Lck, Blk, and Hck, are important in the regulation of growth and differentiation of eukaryotic cells (1). Src activity is regulated by tyrosine phosphorylation at two sites, but with opposing effects. While phosphorylation at Tyr416 in the activation loop of the kinase domain upregulates enzyme activity, phosphorylation at Tyr527 in the carboxy-terminal tail by Csk renders the enzyme less active (2).p56lck (lymphocyte cellular kinase) has been shown to be a pivotal enzyme to both maturation of thymocytes and activation and proliferation of peripheral lymphocytes. The p56lck sequence appeared highly homologous to that of the oncogene p60c-src as did its exon-intron organisation. p56lck does not appear involved in lymphoproliferative diseases, either by overexpression or activating mutations. Nevertheless, its aberrant expression has been reported in some carcinomas (colon, lung and mammary). It was suggested that p56lck could favor metastases by facilitating loss of cell adhesion (3).

To place an order, please [Click HERE](#).