

IFIH1

Reactivity:Human Mouse

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:117kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human IFIH1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b

Synonym:

HLcd; IDDM19; MDA-5; MDA5; MGC133047;

Catalog #:A2203

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:64135

Isotype:IgG

Swiss Prot:Q9BYX4

Purity:Affinity purification

For research use only.

Background:

cts viral dsRNA and two amino-terminal caspase recruitment domains (CARD) that are required for triggering downstream signaling (4-7). Rig-I binds both dsRNA and viral ssRNA that contains a 5'-triphosphate end not seen in host RNA (8,9). Though structurally related, Rig-I and MDA-5 detect a distinct set of viruses (10,11). The CARD domain of the helicases, which is sufficient to generate signaling and IFN production, is recruited to the CARD domain of the MAVS/VISA/Cardif/IPS-1 mitochondrial protein, which triggers activation of NF-B, TBK1/IKK, and IRF-3/IRF-7 (12-15).

To place an order, please [Click HERE](#).