

CKMT2

Reactivity: Human Mouse Rat

Tested applications: WB

Recommended Dilution: WB 1:500 - 1:1000

Calculated MW: 56kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant Protein of human CKMT2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

ag

Synonym:

CHRNA7-2; NACHRA7;

Catalog #: A2206

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 1160

Isotype: IgG

Swiss Prot: P36544

Purity: Affinity purification

For research use only.

Background:

The nicotinic acetylcholine receptors (nAChRs) are members of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. The nAChRs are thought to be hetero-pentamers composed of homologous subunits. The proposed structure for each subunit is a conserved N-terminal extracellular domain followed by three conserved transmembrane domains, a variable cytoplasmic loop, a fourth conserved transmembrane domain, and a short C-terminal extracellular region. The protein encoded by this gene forms a homo-oligomeric channel, displays marked permeability to calcium ions and is a major component of brain nicotinic receptors that are blocked by, and highly sensitive to, alpha-bungarotoxin. Once this receptor binds acetylcholine, it undergoes an extensive change in conformation that affects all subunits and leads to opening of an ion-conducting channel across the plasma membrane. This gene is located in a region identified as a major susceptibility locus for juvenile myoclonic epilepsy and a chromosomal location involved in the genetic transmission of schizophrenia. An evolutionarily recent partial duplication event in this region results in a hybrid containing sequence from this gene and a novel FAM7A gene. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2012]

To place an order, please [Click HERE](#).