

HES1

Reactivity:Human Mouse Rat

Tested applications:WB IHC ICC FC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 ICC 1:50 - 1:200 FC 1:20 - 1:50

Calculated MW:30kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human HES1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

HES1;FLJ20408;HES-1;HHL;HRY;bHLHb39;

Catalog #:A2210

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3280

Isotype:IgG

Swiss Prot:Q14469

Purity:Affinity purification

For research use only.

Background:

The Drosophila Hairy and enhancer of split genes encode basic helix-loop-helix (bHLH) transcriptional repressors that function in the Notch signaling pathway and control segmentation and neural development during embryogenesis. The mammalian homolog of Drosophila Hairy and enhancer of split are the HES gene family members HES1-6, which also encode bHLH transcriptional repressors that regulate myogenesis and neurogenesis. The HES family members form a complex with TLE, the mammalian homolog of groucho, and this interaction is mediated by the carboxy-terminal WRPW motif of the HES proteins. The HES/TLE complex functions by directly binding to DNA instead of interfering with activator proteins. Most HES family members, including HES1 and HES5, preferentially bind to the N box (CACNAG) as opposed to the E box (CANNTG). HES2 binds to both N and E box sites, while HES6 does not bind DNA. Rather, HES6 inhibits HES1 activity, thereby promoting transcription. HES1 and HES2 are expressed in a variety of adult and embryonic tissues. HES3 is expressed exclusively in cerebellar Purkinje cells and HES5 is found solely in the nervous system. HES6 is produced in brain as well as in the limb buds of developing embryos.

To place an order, please [Click HERE](#).