

## ALCAM

**Reactivity:** Human

**Tested applications:** WB

**Recommended Dilution:** WB 1:500 - 1:2000

**Calculated MW:** 100-105kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human ALCAM

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

ALCAM; CD166; MEMD; CD166 antigen;

**Catalog #:** A2218

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 214

**Isotype:** IgG

**Swiss Prot:** Q13740

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes activated leukocyte cell adhesion molecule (ALCAM), also known as CD166 (cluster of differentiation 166), which is a member of a subfamily of immunoglobulin receptors with five immunoglobulin-like domains (VVC2C2C2) in the extracellular domain. This protein binds to T-cell differentiation antigen CD6, and is implicated in the processes of cell adhesion and migration. Multiple alternatively spliced transcript variants encoding different isoforms have been found. [provided by RefSeq, Aug 2011]

**To place an order, please [Click HERE](#).**