

## RNF2

---

**Reactivity:**Human

**Tested applications:**WB

**Recommended Dilution:**WB 1:200 - 1:2000

**Calculated MW:**38kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human RNF2

**Storage Buffer:**

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

fl

**Synonym:**

BAP1; DING; BAP-1; HIPI3; RING2; RING1B;

**Catalog #:**A2232

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6045

**Isotype:**IgG

**Swiss Prot:**Q99496

**Purity:**Affinity purification

For research use only.

**Background:**

Polycomb group (PcG) of proteins form the multiprotein complexes that are important for the transcription repression of various genes involved in development and cell proliferation. The protein encoded by this gene is one of the PcG proteins. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), an ubiquitin-conjugating enzyme, and possess ubiquitin ligase activity.

**To place an order, please [Click HERE](#).**