

HDAC8

Reactivity: Human Mouse Rat

Tested applications: WB IP

Recommended Dilution: WB 1:500 - 1:2000 IP 1:20 - 1:100

Calculated MW: 42kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human HDAC8

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fp

Synonym:

HD8; WTS; RPD3; CDA07; CDLS5; MRXS6; HDACL1;

Catalog #: A2323

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 55869

Isotype: IgG

Swiss Prot: Q9BY41

Purity: Affinity purification

For research use only.

Background:

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).