

## PIK3CG

**Reactivity:**Human Mouse

**Tested applications:**WB IP

**Recommended Dilution:**WB 1:500 - 1:1000 IP 1:20 - 1:100

**Calculated MW:**126kDa

**Observed MW:**Refer to figures

**Immunogen:**

A synthetic peptide of human PIK3CG

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bq

**Synonym:**

PI3K; PIK3; PI3CG; PI3Kgamma; p110gamma; p120-PI3K;

**Catalog #:**A2461

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5294

**Isotype:**IgG

**Swiss Prot:**P48736

**Purity:**Affinity purification

For research use only.

**Background:**

Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene.

**To place an order, please [Click HERE](#).**