

TLR4

Reactivity: Human

Tested applications:WB FC

Recommended Dilution: WB 1:500 - 1:1000 FC 1:20 - 1:50

Calculated MW:96kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human TLR4

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

s

Synonym:

TOLL; CD284; TLR-4; ARMD10

Background:

The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor has been implicated in signal transduction events induced by lipopolysaccharide (LPS) found in most gram-negative bacteria. Mutations in this gene have been associated with differences in LPS responsiveness. Multiple transcript variants encoding different isoforms have been found for this gene.

To place an order, please Click HERE.



Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID:7099

Isotype:IgG

Swiss Prot: 000206

Purity: Affinity purification

For research use only.





