

HNMT

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:33kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human HNMT

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

f

Synonym:

HMT; HNMT-S1; HNMT-S2;

Catalog #:A2520

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3176

Isotype:IgG

Swiss Prot:P50135

Purity:Affinity purification

For research use only.

Background:

In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different proteins have been found for this gene.

To place an order, please [Click HERE](#).