

## PCBP2

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:** 39kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human PCBP2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

HNRPE2; MGC110998; hnRNP-E2;

**Catalog #:** A2531

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 5094

**Isotype:** IgG

**Swiss Prot:** Q15366

**Purity:** Affinity purification

For research use only.

**Background:**

The protein encoded by this gene appears to be multifunctional. Along with PCBP-1 and hnRNP, it is one of the major cellular poly(rC)-binding proteins. The encoded protein contains three K-homologous (KH) domains which may be involved in RNA binding. Together with PCBP-1, this protein also functions as a translational coactivator of poliovirus RNA via a sequence-specific interaction with stem-loop IV of the IRES, promoting poliovirus RNA replication by binding to its 5'-terminal cloverleaf structure. It has also been implicated in translational control of the 15-lipoxygenase mRNA, human papillomavirus type 16 L2 mRNA, and hepatitis A virus RNA. The encoded protein is also suggested to play a part in formation of a sequence-specific alpha-globin mRNP complex which is associated with alpha-globin mRNA stability. This multiexon structural mRNA is thought to be retrotransposed to generate PCBP-1, an intronless gene with functions similar to that of PCBP2. This gene and PCBP-1 have paralogous genes (PCBP3 and PCBP4) which are thought to have arisen as a result of duplication events of entire genes. This gene also has two processed pseudogenes (PCBP2P1 and PCBP2P2). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

**To place an order, please [Click HERE](#).**