

## PSMD3

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**61kDa

**Observed MW:**Refer to Figures

**Immunogen:**

A synthetic peptide of human PSMD3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

S3;P58;RPN3;TSTA2;

**Catalog #:**A2790

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**5709

**Isotype:**IgG

**Swiss Prot:**O43242

**Purity:**Affinity purification

For research use only.

**Background:**

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits.

Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a member of the proteasome subunit S3 family that functions as one of the non-ATPase subunits of the 19S regulator lid. Single nucleotide polymorphisms in this gene are associated with neutrophil count.

*To place an order, please [Click HERE](#).*