www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

IAPP

Reactivity:Human Mouse Rat

Tested applications:WB

Recommended Dilution:WB 1:500 - 1:2000
Calculated MW:10kDa
Observed MW:Refer to Figures
Immunogen:

A synthetic peptide of human IAPP

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

b
Synonym:

DAP; IAP;

Background:

Islet, or insulinoma, amyloid polypeptide is commonly found in pancreatic islets of patients suffering diabetes mellitus type II, or harboring an insulinoma. While the association of amylin with the development of type II diabetes has been known for some time, a direct causative role for amylin has been harder to establish. Studies suggest that amylin, like the related beta-amyloid (Abeta) associated with Alzheimer's disease, can induce apoptotic cell-death in particular cultured cells, an effect that may be relevant to the development of type II diabetes.

To place an order, please Click HERE.



Catalog #:A2816 Antibody Type: Polyclonal Antibody Species:Rabbit Gene ID:3375 Isotype:IgG Swiss Prot:P10997 Purity:Affinity purification

For research use only.



