

CCL3L1

Reactivity: Human

Tested applications: WB

Recommended Dilution: WB 1:200 - 1:500

Calculated MW: 10kDa

Observed MW: Refer to figures

Immunogen:

Recombinant protein of human CCL3L1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

LD78; 464.2; MIP1AP; SCYA3L; G0S19-2; SCYA3L1; D17S1718; LD78BETA; LD78-beta(1-70);

Catalog #: A2852

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 6349

Isotype: IgG

Swiss Prot: P16619

Purity: Affinity purification

For research use only.

Background:

This gene is one of several cytokine genes that are clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins that function in inflammatory and immunoregulatory processes. The protein encoded by this gene binds to several chemokine receptors, including chemokine binding protein 2 and chemokine (C-C motif) receptor 5 (CCR5). CCR5 is a co-receptor for HIV, and binding of this protein to CCR5 inhibits HIV entry. The copy number of this gene varies among individuals, where most individuals have one to six copies, and a minority of individuals have zero or more than six copies. There are conflicting reports about copy number variation of this gene and its correlation to disease susceptibility. This record represents one of two copies that are present on the ALT_REF_LOCI_2 alternate haplotype of the GRCh38 human reference genome assembly. Alternative splicing of this gene results in multiple transcript variants.

To place an order, please [Click HERE](#).