

## CCL13

**Reactivity:** Human

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:100

**Calculated MW:** 11kDa

**Observed MW:** Refer to figures

**Immunogen:**

A synthetic peptide of human CCL13

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

NCC1; CKb10; MCP-4; NCC-1; SCYL1; SCYA13;

**Catalog #:** A2853

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 6357

**Isotype:** IgG

**Swiss Prot:** Q99616

**Purity:** Affinity purification

For research use only.

**Background:**

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during atherosclerosis.

**To place an order, please [Click HERE](#).**