

## CCL17

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:** 11kD

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human CCL17

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

TARC; ABCD-2; SCYA17; A-152E5.3;

**Catalog #:** A2854

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 6361

**Isotype:** IgG

**Swiss Prot:** Q92583

**Purity:** Affinity purification

For research use only.

**Background:**

This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

**To place an order, please [Click HERE](#).**