

## CCL17

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**11kD

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human CCL17

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

TARC; ABCD-2; SCYA17; A-152E5.3;

**Catalog #:**A2854

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6361

**Isotype:**IgG

**Swiss Prot:**Q92583

**Purity:**Affinity purification

For research use only.

**Background:**

This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells.

*To place an order, please [Click HERE](#).*