

## POMC

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:** 29kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human POMC

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

LPH; MSH; NPP; POC; ACTH; CLIP

**Catalog #:** A3038

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 5443

**Isotype:** IgG

**Swiss Prot:** P01189

**Purity:** Affinity purification

For research use only.

**Background:**

This gene encodes a polypeptide hormone precursor that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the polypeptide precursor and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation.

**To place an order, please [Click HERE](#).**