

## NFASC

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:100

**Calculated MW:**150kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human NFASC

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

NF; NRCAML

**Catalog #:**A3053

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**23114

**Isotype:**IgG

**Swiss Prot:**O94856

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes an L1 family immunoglobulin cell adhesion molecule with multiple IGcam and fibronectin domains. The protein functions in neurite outgrowth, neurite fasciculation, and organization of the axon initial segment (AIS) and nodes of Ranvier on axons during early development. Both the AIS and nodes of Ranvier contain high densities of voltage-gated Na<sup>+</sup> (Nav) channels which are clustered by interactions with cytoskeletal and scaffolding proteins including this protein, gliomedin, ankyrin 3 (ankyrin-G), and betaIV spectrin. This protein links the AIS extracellular matrix to the intracellular cytoskeleton. This gene undergoes extensive alternative splicing, and the full-length nature of some variants has not been determined.

**To place an order, please [Click HERE](#).**