

SLC1A4

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW:56kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human SLC1A4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ASCT1; SATT;

Background:

SLC1A4, also known as ASCT1, is a neutral amino acid transporter. Its other name, ASCT1, was given because it mediates obligatory exchange of alanine, serine, cysteine, and threonine (1).

SLC1A4 mediates the efflux of glutamate from the neuron into the synaptic junction via calcium-independent release, as well as mediating the efflux of L-serine from glial cells and its uptake by neurons (2). SLC1A4-mediated transport is shown to involve a symmetrical potassium-independent electroneutral exchange of neutral amino acids and sodium, such that the current activated during transport is carried only by chloride ions (3).

To place an order, please [Click HERE](#).

Catalog #:A3084

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:6509

Isotype:IgG

Swiss Prot:P43007

Purity:Affinity purification

For research use only.