

## TRAF7

**Reactivity:** Human Mouse Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:1000 IHC 1:50 - 1:100

**Calculated MW:** 67kD

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human TRAF7

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

1mg/ml

**Synonym:**

RFWD1; RNF119;

**Catalog #:** A3095

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 84231

**Isotype:** IgG

**Swiss Prot:** Q6Q0C0

**Purity:** Affinity purification

For research use only.

**Background:**

Tumor necrosis factor (TNF; see MIM 191160) receptor-associated factors, such as TRAF7, are signal transducers for members of the TNF receptor superfamily (see MIM 191190). TRAFs are composed of an N-terminal cysteine/histidine-rich region containing zinc RING and/or zinc finger motifs; a coiled-coil (leucine zipper) motif; and a homologous region that defines the TRAF family, the TRAF domain, which is involved in self-association and receptor binding.

**To place an order, please [Click HERE](#).**