

PDE1C

Reactivity:Human Mouse Rat

Tested applications:WB IP

Recommended Dilution:WB 1:500 - 1:2000 IP 1:20 - 1:50

Calculated MW:81kDa

Observed MW:Refer to figures

Immunogen:

A synthetic Peptide of human PDE1C

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

Hcam3; hCam-3; cam-PDE 1C;

Catalog #:A4009

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:5137

Isotype:IgG

Swiss Prot:Q14123

Purity:Affinity purification

For research use only.

Background:

Cyclic nucleotide phosphodiesterases (PDEs) catalyze hydrolysis of the cyclic nucleotides cAMP and cGMP to the corresponding nucleoside 5-prime-monophosphates. Mammalian PDEs have been classified into several families based on their biochemical properties. Members of the PDE1 family, such as PDE1C, are calmodulin (see MIM 114180)-dependent PDEs (CaM-PDEs) that are stimulated by a calcium-calmodulin complex (Repaske et al., 1992

To place an order, please [Click HERE](#).