PTPN7

Reactivity: Human

Tested applications:WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW:41kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human PTPN7

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

0

Synonym:

LPTP; HEPTP; PTPNI; BPTP-4; LC-PTP;

Background:

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Multiple alternatively spliced transcript variants have been found for this gene.

To place an order, please Click HERE.

Polyclonal Antibody

Species: Rabbit

Gene ID:5778

Isotype:IgG

Swiss Prot:P35236

Purity: Affinity purification

For research use only.







