

GALNT4

Reactivity:Human Mouse Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:20 - 1:200

Calculated MW:67kDa

Observed MW:Refer to figures

Immunogen:

A synthetic Peptide of human GALNT4

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GALNACT4; GALNAC-T4;

Catalog #:A4243

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:8693

Isotype:IgG

Swiss Prot:Q8N4A0

Purity:Affinity purification

For research use only.

Background:

This gene encodes a member of the UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GalNAc-T) family of enzymes. GalNAc-Ts initiate mucin-type O-linked glycosylation in the Golgi apparatus by catalyzing the transfer of GalNAc to serine and threonine residues on target proteins. They are characterized by an N-terminal transmembrane domain, a stem region, a luminal catalytic domain containing a GT1 motif and Gal/GalNAc transferase motif, and a C-terminal ricin/lectin-like domain. GalNAc-Ts have different, but overlapping, substrate specificities and patterns of expression. In vitro, the encoded protein can complement other GalNAc-Ts in the complete O-glycosylation of the mucin-1 tandem repeat and can O-glycosylate the P-selectin glycoprotein ligand-1 molecule. The coding region of this gene is contained within a single exon. Fusion transcripts, which combine part of this gene with the 5' exons of the neighboring POC1B (POC1 centriolar protein homolog B) gene, also exist.

To place an order, please [Click HERE](#).