

## RRM2

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

**Calculated MW:**45kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human RRM2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

e

**Synonym:**

R2;RR2M;

**Catalog #:**A5255

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6241

**Isotype:**IgG

**Swiss Prot:**P31350

**Purity:**Affinity purification

For research use only.

**Background:**

Ribonucleotide reductase M2 subunit is one of two subunits that constitute ribonucleotide reductase, the enzyme that catalyzes the conversion of ribonucleotide 5'-diphosphates into 2'-deoxyribonucleotides, a rate-limiting step in the production of 2'-deoxyribonucleoside 5'-diphosphates (dNTP) required for DNA synthesis and repair that is required for DNA synthesis and repair [PMID:20825972, 19250552]. RRM2 is only expressed during the late G1/early S phase, and degraded in late S phase, and the activity of RNR, and therefore DNA synthesis and cell proliferation, is controlled during the cell cycle by the synthesis and degradation of RRM2 subunit.

*To place an order, please [Click HERE](#).*