

## KLC1

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**65kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human KLC1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

KLC; KNS2; KNS2A; RP11-73M18.7; RP11-73M18.8;

**Catalog #:**A5552

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**3831

**Isotype:**IgG

**Swiss Prot:**Q07866

**Purity:**Affinity purification

For research use only.

**Background:**

Conventional kinesin is a tetrameric molecule composed of two heavy chains and two light chains, and transports various cargos along microtubules toward their plus ends. The heavy chains provide the motor activity, while the light chains bind to various cargos. This gene encodes a member of the kinesin light chain family. It associates with kinesin heavy chain through an N-terminal domain, and six tetratricopeptide repeat (TPR) motifs are thought to be involved in binding of cargos such as vesicles, mitochondria, and the Golgi complex. Thus, kinesin light chains function as adapter molecules and not motors per se. Although previously named "kinesin 2", this gene is not a member of the kinesin-2 / kinesin heavy chain subfamily of kinesin motor proteins. Extensive alternative splicing produces isoforms with different C-termini that are proposed to bind to different cargos; however, the full-length nature and/or biological validity of most of these variants have not been determined.

**To place an order, please [Click HERE](#).**