

RAG2

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:59kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human RAG2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

RAG-2;

Background:

This gene encodes a protein that is involved in the initiation of V(D)J recombination during B and T cell development. This protein forms a complex with the product of the adjacent recombination activating gene 1, and this complex can form double-strand breaks by cleaving DNA at conserved recombination signal sequences. The recombination activating gene 1 component is thought to contain most of the catalytic activity, while the N-terminal of the recombination activating gene 2 component is thought to form a six-bladed propeller in the active core that serves as a binding scaffold for the tight association of the complex with DNA. A C-terminal plant homeodomain finger-like motif in this protein is necessary for interactions with chromatin components, specifically with histone H3 that is trimethylated at lysine 4. Mutations in this gene cause Omenn syndrome, a form of severe combined immunodeficiency associated with autoimmune-like symptoms.

To place an order, please [Click HERE](#).

Catalog #:A5626

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:5897

Isotype:IgG

Swiss Prot:P55895

Purity:Affinity purification

For research use only.