

FH

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:10 - 1:100

Calculated MW:54kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human FH

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

MCL; LRCC; HLRCC; MCUL1;

Catalog #:A5688

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2271

Isotype:IgG

Swiss Prot:P07954

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Krebs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form is targeted to the mitochondrion, where the removal of the extension generates the same form as in the cytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutations in this gene can cause fumarase deficiency and lead to progressive encephalopathy.

To place an order, please [Click HERE](#).