

## GPD1

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**37kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human GPD1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

GPD-C; HTGTI; GPDH-C;

**Catalog #:**A5715

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**2819

**Isotype:**IgG

**Swiss Prot:**P21695

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes a member of the NAD-dependent glycerol-3-phosphate dehydrogenase family. The encoded protein plays a critical role in carbohydrate and lipid metabolism by catalyzing the reversible conversion of dihydroxyacetone phosphate (DHAP) and reduced nicotinic adenine dinucleotide (NADH) to glycerol-3-phosphate (G3P) and NAD<sup>+</sup>. The encoded cytosolic protein and mitochondrial glycerol-3-phosphate dehydrogenase also form a glycerol phosphate shuttle that facilitates the transfer of reducing equivalents from the cytosol to mitochondria. Mutations in this gene are a cause of transient infantile hypertriglyceridemia. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

*To place an order, please [Click HERE](#).*