

PSMA5

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW: 26kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human PSMA5

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

PSC5; ZETA;

Catalog #: A5752

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 5686

Isotype: IgG

Swiss Prot: P28066

Purity: Affinity purification

For research use only.

Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Multiple alternatively spliced transcript variants encoding two distinct isoforms have been found for this gene.

To place an order, please [Click HERE](#).