

## TDG

---

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**46kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human TDG

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

ef

**Synonym:**

hTDG;

**Catalog #:**A5756

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6996

**Isotype:**IgG

**Swiss Prot:**Q13569

**Purity:**Affinity purification

For research use only.

**Background:**

The protein encoded by this gene belongs to the TDG/mug DNA glycosylase family. Thymine-DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired thymine. With lower activity, this enzyme also removes thymine from C/T and T/T mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. This enzyme plays a central role in cellular defense against genetic mutation caused by the spontaneous deamination of 5-methylcytosine and cytosine. This gene may have a pseudogene in the p arm of chromosome 12.

*To place an order, please [Click HERE](#).*