

## MRPS30

**Reactivity:**Human Mouse

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

**Calculated MW:**50kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human MRPS30

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

bj

**Synonym:**

PAP; PDCD9; S30mt; MRP-S30;

**Catalog #:**A5841

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**10884

**Isotype:**IgG

**Swiss Prot:**Q9NP92

**Purity:**Affinity purification

For research use only.

**Background:**

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that is similar to the chicken pro-apoptotic protein p52. Transcript variants using alternative promoters or polyA sites have been mentioned in the literature but the complete description of these sequences is not available.

**To place an order, please [Click HERE](#).**