

ILF2

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:43kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human ILF2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

NF45; PRO3063;

Catalog #:A5882

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:3608

Isotype:IgG

Swiss Prot:Q12905

Purity:Affinity purification

For research use only.

Background:

The protein encoded by this gene is the 45 kDa component of nuclear factor of activated T-cells (NFAT), a heterodimer of 45 kDa and 90 kDa proteins. NFAT is a transcription factor required for T-cell expression of the interleukin 2 gene. It also binds RNA and is an essential component for encapsidation and protein priming of hepatitis B viral polymerase. The complex has been shown to repair DNA breaks by nonhomologous end joining and can also negatively regulate the microRNA processing pathway. Alternative splicing results in multiple transcript variants. Related pseudogenes have been found on chromosomes 3 and 14.

To place an order, please [Click HERE](#).