www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

HTATSF1

Reactivity: Human Mouse

Tested applications: WB IHC

Recommended Dilution:WB 1:1000 - 1:2000 IHC 1:50 - 1:100

Calculated MW:86kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human HTATSF1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

Synonym:

TATSF1; TAT-SF1; dJ196E23.2;

Background:

The protein encoded by this gene functions as a cofactor for the stimulation of transcriptional elongation by HIV-1 Tat, which binds to the HIV-1 promoter through Tat-TAR interaction. This protein may also serve as a dual-function factor to couple transcription and splicing and to facilitate their reciprocal activation. Alternatively spliced transcript variants have been found for this gene.

To place an order, please Click HERE.

Catalog #:A5977

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:27336

Isotype:IgG

Swiss Prot: 043719

Purity: Affinity purification

For research use only.





