

PTRF

Reactivity:Human

Tested applications:WB IHC

Recommended Dilution:WB 1:1000 - 1:2000 IHC 1:50 - 1:100

Calculated MW:43kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human PTRF

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

j

Synonym:

CGL4; CAVIN; CAVIN1; FKSG13; cavin-1;

Catalog #:A6007

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:284119

Isotype:IgG

Swiss Prot:Q6NZI2

Purity:Affinity purification

For research use only.

Background:

This gene encodes a protein that enables the dissociation of paused ternary polymerase I transcription complexes from the 3' end of pre-rRNA transcripts. This protein regulates rRNA transcription by promoting the dissociation of transcription complexes and the reinitiation of polymerase I on nascent rRNA transcripts. This protein also localizes to caveolae at the plasma membrane and is thought to play a critical role in the formation of caveolae and the stabilization of caveolins. This protein translocates from caveolae to the cytoplasm after insulin stimulation. Caveolae contain truncated forms of this protein and may be the site of phosphorylation-dependent proteolysis. This protein is also thought to modify lipid metabolism and insulin-regulated gene expression. Mutations in this gene result in a disorder characterized by generalized lipodystrophy and muscular dystrophy.

To place an order, please [Click HERE](#).