

## NXF1

**Reactivity:** Human Mouse Rat Hamster

**Tested applications:** WB IP

**Recommended Dilution:** WB 1:500 - 1:1000 IP 1:20 - 1:50

**Calculated MW:** 70kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human NXF1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

TAP; MEX67;

**Catalog #:** A6040

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 10482

**Isotype:** IgG

**Swiss Prot:** Q9UBU9

**Purity:** Affinity purification

For research use only.

**Background:**

This gene is one member of a family of nuclear RNA export factor genes. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. Alternative splicing seems to be a common mechanism in this gene family. The encoded protein of this gene shuttles between the nucleus and the cytoplasm and binds in vivo to poly(A)<sup>+</sup> RNA. It is the vertebrate homologue of the yeast protein Mex67p. The encoded protein overcomes the mRNA export block caused by the presence of saturating amounts of CTE (constitutive transport element) RNA of type D retroviruses. Alternative splicing results in multiple transcript variants.

**To place an order, please [Click HERE](#).**