www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## PRPF8



Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:1000 IHC 1:50 - 1:100

Calculated MW:273kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human PRPF8

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

Synonym:

PRP8; RP13; HPRP8; PRPC8; SNRNP220;

Background:

Pre-mRNA splicing occurs in 2 sequential transesterification steps. The protein encoded by this gene is a component of both U2- and U12-dependent spliceosomes, and found to be essential for the catalytic step II in pre-mRNA splicing process. It contains several WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to yeast Prp8 protein. This gene is a candidate gene for autosomal dominant retinitis pigmentosa.

To place an order, please Click HERE.

Catalog #:A6053

**Antibody Type:** 

Polyclonal Antibody

Species:Rabbit

Gene ID:10594

Isotype:IgG

Swiss Prot:Q6P2Q9

Purity: Affinity purification

For research use only.



