www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## SRSF3

Reactivity: Human Mouse Rat Hamster

Tested applications: WB ICC IP

Recommended Dilution: WB 1:500 - 1:1000 ICC 1:50 - 1:200 IP 1:20 - 1:50

Calculated MW:19kDa

Observed MW:Refer to Figures

Immunogen:

A synthetic peptide of human SRSF3

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Concentration:

Synonym:

SFRS3; SRp20;

## Background:

The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants, one protein-coding and the other non-coding, have been found for this gene.

To place an order, please Click HERE.



Catalog #:A6067

**Antibody Type:** 

Polyclonal Antibody

Species: Rabbit

Gene ID:6428 Isotype:IgG

Swiss Prot:P84103

Purity: Affinity purification

For research use only.





