

## PTBP1

**Reactivity:** Human Rat

**Tested applications:** WB IHC

**Recommended Dilution:** WB 1:500 - 1:1000 IHC 1:100 - 1:200

**Calculated MW:** 57kDa

**Observed MW:** Refer to Figures

**Immunogen:**

A synthetic peptide of human PTBP1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

j

**Synonym:**

PTB; PTB2; PTB3; PTB4; pPTB; HNRPI; PTB-1; PTB-T; HNRNP1; HNRNP-I;

**Catalog #:** A6107

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 5725

**Isotype:** IgG

**Swiss Prot:** P26599

**Purity:** Affinity purification

For research use only.

**Background:**

This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA-binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has four repeats of quasi-RNA recognition motif (RRM) domains that bind RNAs. This protein binds to the intronic polypyrimidine tracts that requires pre-mRNA splicing and acts via the protein degradation ubiquitin-proteasome pathway. It may also promote the binding of U2 snRNP to pre-mRNAs. This protein is localized in the nucleoplasm and it is also detected in the perinucleolar structure. Alternatively spliced transcript variants encoding different isoforms have been described.

**To place an order, please [Click HERE](#).**