

## SF3B3

**Reactivity:**Human Mouse Rat

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**136kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human SF3B3

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:**

f

**Synonym:**

RSE1; SAP130; SF3b130; STAF130;

**Catalog #:**A6146

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**23450

**Isotype:**IgG

**Swiss Prot:**Q15393

**Purity:**Affinity purification

For research use only.

**Background:**

This gene encodes subunit 3 of the splicing factor 3b protein complex. Splicing factor 3b, together with splicing factor 3a and a 12S RNA unit, forms the U2 small nuclear ribonucleoproteins complex (U2 snRNP). The splicing factor 3b/3a complex binds pre-mRNA upstream of the intron's branch site in a sequence independent manner and may anchor the U2 snRNP to the pre-mRNA. Splicing factor 3b is also a component of the minor U12-type spliceosome. Subunit 3 has also been identified as a component of the STAGA (SPT3-TAF(II)31-GCN5L acetylase) transcription coactivator-HAT (histone acetyltransferase) complex, and the TFTC (TATA-binding-protein-free TAF(II)-containing complex). These complexes may function in chromatin modification, transcription, splicing, and DNA repair.

*To place an order, please [Click HERE](#).*