

## PHLDA2

**Reactivity:** Human

**Tested applications:** WB IHC IF

**Recommended Dilution:** WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:20 - 1:100

**Calculated MW:** 17kDa

**Observed MW:** Refer to Figures

**Immunogen:**

Recombinant protein of human PHLDA2

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

IPL; BRW1C; BWR1C; HLDA2; TSSC3;

**Catalog #:** A6244

**Antibody Type:**

Polyclonal Antibody

**Species:** Rabbit

**Gene ID:** 7262

**Isotype:** IgG

**Swiss Prot:** Q53GA4

**Purity:** Affinity purification

For research use only.

**Background:**

This gene is located in a cluster of imprinted genes on chromosome 11p15.5, which is considered to be an important tumor suppressor gene region. Alterations in this region may be associated with the Beckwith-Wiedemann syndrome, Wilms tumor, rhabdomyosarcoma, adrenocortical carcinoma, and lung, ovarian, and breast cancer. This gene has been shown to be imprinted, with preferential expression from the maternal allele in placenta and liver.

*To place an order, please [Click HERE](#).*