

GCK

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

Calculated MW:52kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human GCK

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

GK; GLK; HK4; HHF3; HKIV; HXKP; LGLK; MODY2; FGQTL3;

Background:

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. Alternative splicing of this gene results in three tissue-specific forms of glucokinase, one found in pancreatic islet beta cells and two found in liver. The protein localizes to the outer membrane of mitochondria. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. Mutations in this gene have been associated with non-insulin dependent diabetes mellitus (NIDDM), maturity onset diabetes of the young, type 2 (MODY2) and persistent hyperinsulinemic hypoglycemia of infancy (PHHI).

To place an order, please [Click HERE](#).

Catalog #:A6293

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:2645

Isotype:IgG

Swiss Prot:P35557

Purity:Affinity purification

For research use only.