

## XCL1

---

**Reactivity:**Human

**Tested applications:**WB IHC IF

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:100

**Calculated MW:**12kDa

**Observed MW:**Refer to Figures

**Immunogen:**

Recombinant protein of human XCL1

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

LTN; ATAC; LPTN; SCM1; SCM-1; SCM1A; SCYC1; SCM-1a;

**Background:**

This antimicrobial gene encodes a member of the chemokine superfamily. Chemokines function in inflammatory and immunological responses, inducing leukocyte migration and activation. The encoded protein is a member of the C-chemokine subfamily, retaining only two of four cysteines conserved in other chemokines, and is thought to be specifically chemotactic for T cells. This gene and a closely related family member are located on the long arm of chromosome 1.

*To place an order, please [Click HERE](#).*

**Catalog #:**A6408

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**6375

**Isotype:**IgG

**Swiss Prot:**P47992

**Purity:**Affinity purification

For research use only.