

NR2C2

Reactivity:Human Rat

Tested applications:WB IHC

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW:65kDa

Observed MW:Refer to Figures

Immunogen:

Recombinant protein of human NR2C2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

TR4; TAK1;

Background:

This gene encodes a protein that belongs to the nuclear hormone receptor family. Members of this family act as ligand-activated transcription factors and function in many biological processes such as development, cellular differentiation and homeostasis. The activated receptor/ligand complex is translocated to the nucleus where it binds to hormone response elements of target genes. The protein encoded by this gene plays a role in protecting cells from oxidative stress and damage induced by ionizing radiation. The lack of a similar gene in mouse results in growth retardation, severe spinal curvature, subfertility, premature aging, and prostatic intraepithelial neoplasia (PIN) development. Alternative splicing results in multiple transcript variants encoding different isoforms.

To place an order, please [Click HERE](#).

Catalog #:A6422

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:7182

Isotype:IgG

Swiss Prot:P49116

Purity:Affinity purification

For research use only.