

CCT4

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:57kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human CCT4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

SRB; Cctd; CCT-DELTA;

Catalog #:A6548

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:10575

Isotype:IgG

Swiss Prot:P50991

Purity:Affinity purification

For research use only.

Background:

The chaperonin containing TCP1 (MIM 186980) complex (CCT), also called the TCP1 ring complex, consists of 2 back-to-back rings, each containing 8 unique but homologous subunits, such as CCT4. CCT assists the folding of newly translated polypeptide substrates through multiple rounds of ATP-driven release and rebinding of partially folded intermediate forms. Substrates of CCT include the cytoskeletal proteins actin (see MIM 102560) and tubulin (see MIM 191130), as well as alpha-transducin (MIM 139330) (Won et al., 1998)

To place an order, please [Click HERE](#).