

NSFL1C

Reactivity:Human Mouse Rat

Tested applications:WB IHC IF

Recommended Dilution:WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW:40kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human NSFL1C

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

P47; UBX1; UBXD10; UBXN2C; dJ776F14.1;

Background:

N-ethylmaleimide-sensitive factor (NSF) and valosin-containing protein (p97) are two ATPases known to be involved in transport vesicle/target membrane fusion and fusions between membrane compartments. A trimer of the protein encoded by this gene binds a hexamer of cytosolic p97 and is required for p97-mediated regrowth of Golgi cisternae from mitotic Golgi fragments. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 8.

To place an order, please [Click HERE](#).

Catalog #:A6677

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:55968

Isotype:IgG

Swiss Prot:Q9UNZ2

Purity:Affinity purification

For research use only.