

## TAF1C

---

**Reactivity:**Human Mouse

**Tested applications:**WB IHC

**Recommended Dilution:**WB 1:500 - 1:2000 IHC 1:50 - 1:200

**Calculated MW:**95kDa

**Observed MW:**Refer to figures

**Immunogen:**

Recombinant protein of human TAF1C

**Storage Buffer:**

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Synonym:**

SL1; TAFI95; TAFI110; MGC:39976;

**Background:**

Initiation of transcription by RNA polymerase I requires the formation of a complex composed of the TATA-binding protein (TBP) and three TBP-associated factors (TAFs) specific for RNA polymerase I. This complex, known as SL1, binds to the core promoter of ribosomal RNA genes to position the polymerase properly and acts as a channel for regulatory signals. This gene encodes the largest SL1-specific TAF. Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

*To place an order, please [Click HERE](#).*

**Catalog #:**A6759

**Antibody Type:**

Polyclonal Antibody

**Species:**Rabbit

**Gene ID:**9013

**Isotype:**IgG

**Swiss Prot:**Q15572

**Purity:**Affinity purification

For research use only.