

FANCL

Reactivity:Human Mouse

Tested applications:WB IHC

Recommended Dilution:WB 1:200 - 1:2000 IHC 1:20 - 1:200

Calculated MW:43kDa

Observed MW:Refer to figures

Immunogen:

Recombinant protein of human FANCL

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

POG; PHF9; FAAP43;

Catalog #:A6812

Antibody Type:

Polyclonal Antibody

Species:Rabbit

Gene ID:55120

Isotype:IgG

Swiss Prot:Q9NW38

Purity:Affinity purification

For research use only.

Background:

The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity; they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group L. Alternative splicing results in two transcript variants encoding different isoforms.

To place an order, please [Click HERE](#).